

**VYVGART Hytrulo®**  
(efgartigimod alfa and hyaluronidase-qvfc)

## GET STARTED WITH YOUR IN-OFFICE **SUBCUTANEOUS INJECTION**

VYVGART Hytrulo offers  
a ~30 to 90-second  
subcutaneous injection\*

\*For at least 30 minutes after your injection, a healthcare professional will monitor you for allergic reactions.  
AChR=acetylcholine receptor.

### **What is VYVGART HYTRULO® (efgartigimod alfa and hyaluronidase-qvfc)?**

VYVGART HYTRULO is a prescription medicine used to treat adults with generalized myasthenia gravis (gMG) who are anti-acetylcholine receptor (AChR) antibody positive.

It is not known if VYVGART HYTRULO is safe and effective in children.

### **SELECT IMPORTANT SAFETY INFORMATION**

Do not take VYVGART HYTRULO if you are allergic to efgartigimod alfa, hyaluronidase, or any of the ingredients in VYVGART HYTRULO. VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.


**Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information for VYVGART HYTRULO](#).**



## A treatment plan that's designed for me

**VYVGART Hytrulo is given in treatment cycles with a personalized break between each cycle**

Each treatment cycle is 4 treatments, 1 given each week.



You'll work with your doctor to personalize your breaks between cycles based on your symptoms and side effects



When your symptoms start to return, talk to your doctor to help them decide when you need another cycle

The safety of starting the next treatment cycle sooner than 4 weeks from the end of the previous treatment cycle has not been established. If you miss a treatment, VYVGART Hytrulo may be given up to 3 days after the missed treatment. After that, continue with the normal time between treatments (dosing schedule). Talk to your doctor if you have any questions about your dosing schedule.

Personalized breaks may help you get the most out of your treatment plan. Learn about real patients' breaks between cycles on [VYVGARTDosing.com](https://www.vyvgartdosing.com).

## SELECT IMPORTANT SAFETY INFORMATION

Before taking VYVGART HYTRULO, tell your healthcare provider about all of your medical conditions, including if you: have an infection or fever; have recently received or are scheduled to receive any vaccinations; have any history of allergic reactions; have kidney (renal) problems. Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

**Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information for VYVGART HYTRULO](#).**

**The day before your subcutaneous injection, you may want to:**

- ✓ Prepare for your visit. You may need to find a ride if you can't drive yourself, and you should plan to get to your appointment at least 15 minutes early
- ✓ Plan to wear clothing that's loose around the waist on your injection day
- ✓ Write down a list of your current medications to bring to your appointment

**Keep in mind: how to pass the time**

VYVGART Hytrulo takes about 30 to 90 seconds to inject, but your healthcare provider will monitor you for allergic reactions for at least 30 minutes after your injection. You may want to bring something to do while you wait.

**Book****Crossword****TV show****Podcast****Game****SELECT IMPORTANT SAFETY INFORMATION**

Before taking VYVGART HYTRULO, tell your healthcare provider about all of your medical conditions, including if you: are pregnant or plan to become pregnant; are breastfeeding or plan to breastfeed. It is not known whether VYVGART HYTRULO will harm your unborn baby. It is not known if VYVGART HYTRULO passes into your breast milk. Pregnancy Exposure Registry. There is a pregnancy exposure registry for women who use VYVGART HYTRULO during pregnancy. The purpose of this registry is to collect information about your health and your baby. Your healthcare provider can enroll you in this registry. You may also enroll yourself or get more information about the registry by calling 1-855-272-6524 or going to [VYVGARTPregnancy.com](https://www.vyvgartpregnancy.com).

**Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information for VYVGART HYTRULO](#).**

## The 5 steps of your VYVGART Hytrulo subcutaneous injection

### 01 | Check in

Once you get to your treatment location, check in with the staff and confirm your insurance information.

### 02 | Get ready

An office staff member will help get you ready for your subcutaneous injection. VYVGART Hytrulo may also be prepared for injection at this time.

### 03 | Receive your injection

A healthcare professional will inject VYVGART Hytrulo subcutaneously into an area of pinched skin on your abdomen over ~30 to 90 seconds. A different area of your abdomen will be used for subsequent injections of VYVGART Hytrulo.

### 04 | Stay for monitoring

You will be monitored for injection site allergic reactions and other side effects for at least 30 minutes after your injection. Consider bringing something to pass the time.

### 05 | Check out

Before you leave, make sure to confirm or schedule your next appointment. If this is the last injection of your treatment cycle, you may want to have a plan in place with your doctor for symptom tracking.



Not actual size

After starting VYVGART Hytrulo, track your symptoms each week to help your doctor determine your personalized breaks.



Scan the QR code or visit [gMGSymptomTracker.com](https://gMGSymptomTracker.com) to find a symptom tracking tool that may help with your VYVGART Hytrulo journey.

Are you enrolled in the VYVGART patient support program?

My VYVGART® Path

With My VYVGART Path, a Nurse Case Manager is waiting to help you throughout your VYVGART Hytrulo treatment journey. **Talk with your doctor** and ask to be enrolled in My VYVGART Path, **or visit [MyVYVGARTPath.com](https://MyVYVGARTPath.com) to download the form.**

Already enrolled? Reach out to a Nurse Case Manager for support at **1-833-MY-PATH-1** (1-833-697-2841).

### SELECT IMPORTANT SAFETY INFORMATION

VYVGART HYTRULO can cause side effects which can be serious, including infection. VYVGART HYTRULO may increase the risk of infection. If you have an active infection, your healthcare provider should delay your treatment with VYVGART HYTRULO until your infection is gone. Tell your healthcare provider right away if you get any of the following signs and symptoms of an infection: fever; chills; frequent and painful urination; cough; pain and blockage of nasal passages; wheezing; shortness of breath; sore throat; excess phlegm; nasal discharge.

**Please see Important Safety Information throughout. Please see full [Prescribing and Patient Information for VYVGART HYTRULO](#).**

Do not take VYVGART HYTRULO if you are allergic to efgartigimod alfa, hyaluronidase, or any of the ingredients in VYVGART HYTRULO. VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

**Before taking VYVGART HYTRULO, tell your healthcare provider about all of your medical conditions, including if you:**

- have an infection or fever.
- have recently received or are scheduled to receive any vaccinations.
- have any history of allergic reactions.
- have kidney (renal) problems.
- are pregnant or plan to become pregnant. It is not known whether VYVGART HYTRULO will harm your unborn baby.
  - Pregnancy Exposure Registry. There is a pregnancy exposure registry for women who use VYVGART HYTRULO during pregnancy. The purpose of this registry is to collect information about your health and your baby. Your healthcare provider can enroll you in this registry. You may also enroll yourself or get more information about the registry by calling 1-855-272-6524 or going to [VYVGARTPregnancy.com](http://VYVGARTPregnancy.com)
- are breastfeeding or plan to breastfeed. It is not known if VYVGART HYTRULO passes into your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

**VYVGART HYTRULO can cause side effects which can be serious, including:**

- **Infection.** VYVGART HYTRULO may increase the risk of infection. If you have an active infection, your healthcare provider should delay your treatment with VYVGART HYTRULO until your infection is gone. Tell your healthcare provider right away if you get any of the following signs and symptoms of an infection:

- fever	- wheezing
- chills	- shortness of breath
- frequent and painful urination	- sore throat
- cough	- excess phlegm
- pain and blockage of nasal passages	- nasal discharge

- **Allergic reactions (hypersensitivity reactions).** VYVGART HYTRULO can cause allergic reactions that can be severe. These reactions can happen during, shortly after, or weeks after your VYVGART HYTRULO injection. Tell your healthcare provider or get emergency help right away if you have any of the following symptoms of an allergic reaction:
  - rash
  - swelling of the face, lips, throat, or tongue
  - shortness of breath
  - hives
  - trouble breathing
  - low blood pressure
  - fainting
- **Infusion or injection-related reactions.** VYVGART HYTRULO can cause infusion or injection-related reactions. These reactions can happen during or shortly after your VYVGART HYTRULO injection. Tell your healthcare provider if you have any of the following symptoms of an infusion or injection-related reaction:
  - high blood pressure
  - chills
  - shivering
  - chest, stomach, or back pain

**The most common side effects of VYVGART HYTRULO include respiratory tract infection, headache, urinary tract infection, and injection site reactions.**

These are not all the possible side effects of VYVGART HYTRULO. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### **What is VYVGART HYTRULO® (efgartigimod alfa and hyaluronidase-qvfc)?**

VYVGART HYTRULO is a prescription medicine used to treat adults with generalized myasthenia gravis (gMG) who are anti-acetylcholine receptor (AChR) antibody positive.

It is not known if VYVGART HYTRULO is safe and effective in children.

**Please see full [Prescribing and Patient Information](#) for VYVGART HYTRULO.**

### **Dosage form and strength:**

VYVGART Hytrulo is available as a single-dose subcutaneous injection containing 180 mg/mL of efgartigimod alfa and 2,000 U/mL of hyaluronidase per vial.