



# VYVGART Infusions

---


Information on the infusion process:  
before, during, and after




## VYVGART dosing—make the most of a treatment plan that's designed for you

**VYVGART is given in treatment cycles with a personalized break between each cycle**


Each treatment cycle is 4 treatments, 1 given each week.



Your first  
treatment cycle



Your personalized break  
is your time off between  
receiving treatment cycles



Your next  
treatment cycle

### How personalized breaks may help you get the most out of your treatment plan



You'll work with your doctor to personalize your breaks between cycles based on your symptoms and side effects.



When your symptoms start to return, talk to your doctor to help them decide when you need another cycle.

For example: Your doctor may start you with a 4-week break after each of your first 3 treatment cycles. Your doctor will continue to adjust your time between additional cycles, based on your symptoms and side effects.

The safety of starting the next treatment cycle sooner than 4 weeks from the last treatment of the previous treatment cycle has not been established. If you miss a scheduled treatment, VYVGART may be given up to 3 days after the scheduled treatment day.

### What is VYVGART® (efgartigimod alfa-fcab)?

VYVGART is a prescription medicine used to treat a condition called generalized myasthenia gravis, which causes muscles to tire and weaken easily throughout the body, in adults who are positive for antibodies directed toward a protein called acetylcholine receptor (anti-AChR antibody positive).

### SELECT IMPORTANT SAFETY INFORMATION

Do not use VYVGART if you have a serious allergy to efgartigimod alfa or any of the other ingredients in VYVGART. VYVGART can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

**Please see Important Safety Information throughout. Please see full [Prescribing Information](#) for VYVGART.**

# YOUR STEP-BY-STEP INFUSION GUIDE

**When you begin any treatment, it's good to be prepared.** Reviewing this checklist may help you plan for your visits to your infusion location.



**Check the boxes** as you prepare for your infusion

## THE DAY BEFORE YOUR INFUSION, WE RECOMMEND THAT YOU:



**Drink plenty of water** and eat how you normally would



**Try** to get a full night of sleep



**Make a list** of your current medications to bring to your appointment



**Plan** how you are getting to the infusion location. You may need to find a ride if you cannot drive yourself

## SELECT IMPORTANT SAFETY INFORMATION

VYVGART may cause serious side effects, including infection. VYVGART may increase the risk of infection. The most common infections were urinary tract and respiratory tract infections. Signs or symptoms of an infection may include fever, chills, frequent and/or painful urination, cough, pain and blockage of nasal passages/sinus, wheezing, shortness of breath, fatigue, sore throat, excess phlegm, nasal discharge, back pain, and/or chest pain.

**Please see Important Safety Information throughout. Please see full [Prescribing Information](#) for VYVGART.**

### ON THE DAY OF YOUR INFUSION, WE RECOMMEND THAT YOU:

- ☐ **Wear** comfortable clothes and bring layers in case it's chilly
- ☐ **Drink** plenty of water
- ☐ **Eat** how you normally would
- ☐ Get to your appointment at least **15 minutes early**

### DURING EACH INFUSION, A HEALTHCARE PROFESSIONAL WILL:



**Prepare  
VYVGART**  
for infusion



**Check your vital  
signs**, including  
your pulse, blood  
pressure, and  
temperature



**Find a vein**  
in your arm and  
put in the  
intravenous  
(IV) line



**Monitor you**  
during and after  
your infusion

You may want something to **pass the time** during the infusion process. Consider bringing a:



**Book**



**Crossword**



**TV show**



**Podcast**



**Game**

### IMMEDIATELY AFTER YOUR INFUSION APPOINTMENT:

- ☐ **Your infusion may take approximately 1 hour.** During this time and for an hour after your infusion, a doctor or nurse will monitor you for any reactions
- ☐ **You will need 4 infusions total to complete the cycle,** so make sure to book your next infusion if you have not done so already

### AFTER YOUR INFUSION:

- ☐ **Carefully track your symptoms.** To do this, you may want to regularly complete a Myasthenia Gravis Activities of Daily Living (MG-ADL) assessment. A My VYVGART Path Nurse Case Manager can teach you how to do this
- ☐ **Talk with your doctor** about how you are feeling
- ☐ **Tell your healthcare provider** immediately about any allergic reactions, such as rashes, swelling under the skin, and shortness of breath

### Still have questions about your VYVGART treatment?

Speak with your doctor if you have any questions or if you want more information about VYVGART.

If you are enrolled in My VYVGART Path, reach out to a Nurse Case Manager for support at **1-833-MY-PATH-1** (1-833-697-2841).

### Not yet enrolled in My VYVGART Path?

**ASK YOUR  
DOCTOR  
TO SIGN YOU UP**



**With My VYVGART Path, a Nurse Case Manager is waiting to help you through your VYVGART treatment journey.**

## IMPORTANT SAFETY INFORMATION

Do not use VYVGART if you have a serious allergy to efgartigimod alfa or any of the other ingredients in VYVGART. VYVGART can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

### VYVGART may cause serious side effects, including:

- **Infection.** VYVGART may increase the risk of infection. The most common infections were urinary tract and respiratory tract infections. Signs or symptoms of an infection may include fever, chills, frequent and/or painful urination, cough, pain and blockage of nasal passages/sinus, wheezing, shortness of breath, fatigue, sore throat, excess phlegm, nasal discharge, back pain, and/or chest pain.
- **Allergic Reactions (hypersensitivity reactions).** VYVGART can cause allergic reactions such as rashes, swelling under the skin, and shortness of breath. Serious allergic reactions, such as trouble breathing and decrease in blood pressure leading to fainting have been reported with VYVGART.
- **Infusion-Related Reactions.** VYVGART can cause infusion-related reactions. The most frequent symptoms and signs reported with VYVGART were high blood pressure, chills, shivering, and chest, abdominal, and back pain.

Tell your doctor if you have signs or symptoms of an infection, allergic reaction, or infusion-related reaction. These can happen while you are receiving your VYVGART treatment or afterward. Your doctor may need to pause or stop your treatment. Contact your doctor immediately if you have signs or symptoms of a serious allergic reaction.

### Before taking VYVGART, tell your doctor if you:

- take any medicines, including prescription and non-prescription medicines, supplements, or herbal medicines,
- have received or are scheduled to receive a vaccine (immunization), or
- have any allergies or medical conditions, including if you are pregnant or planning to become pregnant, or are breastfeeding.

### What are the common side effects of VYVGART?

The most common side effects of VYVGART are respiratory tract infection, headache, and urinary tract infection.

These are not all the possible side effects of VYVGART. Call your doctor for medical advice about side effects. You may report side effects to the US Food and Drug Administration at 1-800-FDA-1088.

### What is VYVGART® (efgartigimod alfa-fcab)?

VYVGART is a prescription medicine used to treat a condition called generalized myasthenia gravis, which causes muscles to tire and weaken easily throughout the body, in adults who are positive for antibodies directed toward a protein called acetylcholine receptor (anti-AChR antibody positive).

### Please see full Prescribing Information for VYVGART.

### Dosage form and strength:

VYVGART is available as a single-dose injection for intravenous use containing 400 mg/20 mL of efgartigimod alfa-fcab per vial.